Ca-P Bol[™]

Dietetic calcium and phosphorus bolus for reduction of the risk of milk fever

Ca-P Bol is a fast releasing bolus, containing a combination of rapidly and easily absorbable calcium, phosphorus and magnesium, which together effectively reduces the risk of milk fever and phosphorus deficiency after calving.

Ca-P Bol contains:

- Calcium chloride optimizes rapidly the calcium blood level
- Monocalcium phosphate and calcium propionate supports calcium and phosphorus blood level for several hours
- Magnesium contributes to a higher level of mobilization of calcium in the blood after calving

Packaging:

- 4 pcs. per zip lock bag
- 20 pcs. per case
- 6 cases per box

Ca-P Bol are covered by EU patent

Benefits and effect:

- Dissolves in the rumen within 25 minutes
- Reduces the risk of milk fever
- Provides 40 grams of easy absorbable calcium and 6 grams of phosphorus
- Reduces the prevalence of downer cows
- Provides sufficient calcium and phosphorus in the blood after calving
- Minimizes the risk of mistakes the cow gets the intended amount of calcium
- Easy to administer with a bolus applicator

GENERALFARM veterinaria e zootecnia

GENERALFARM s.r.l. - Via Umbria, 32 - 42122 Reggio Emilia - Tel. 0522 514251 - Fax 0522 514376 www.generalfarm.it - e-mail: info@generalfarm.it

Application:

 $\mathbf{1}^{^{\mathrm{st}}}$ bolus: administer immediately after calving

D BOI

Ca-P Bol is

covered by

EU patent

2nd bolus: administer 10-14 hours after calving

High-risk cows may be supplemented with additional one Ca-P Bol 24 hours before calving and one 24 hours after calving.

Follow-up on calcium infusion:

st bolus: 2 hours after calcium infusion
2nd bolus: 12 hours after calcium infusion

Consumption:

• 2-4 boluses per cow in the period around calving

